



AnimaXx Premium

# TOP DRY - Litter Conditioner

The new generation of litter conditioner

## Drier litter and healthier environment for animal welfare

### PROPERTIES:

- ✓ Good water absorption capacity 150-170%
- ✓ Phosphorus free, biodegradable
- ✓ Pleasant smell from menthol
- ✓ Makes cleaning easier
- ✓ Keep fly, mosquitos and moth away

### USAGE, DOSAGE:

#### CATLE:

Stall: 400 g per cow, 3x per week  
 Box: 180 g per cow per day  
 For calves: 80 g/m<sup>2</sup>  
 Dry off hooves (in tray)

#### SOW, PIGLET BATHING:

Sow: 65 g per day behind the sow and in farrowing area. For easier cleaning!

Farrowing: Spread Top Dry on the pen and on heat mats before farrowing. Top Dry is especially designed for newborn piglets drying at birth.

Gilt: 100 g per day on straw or sand.

### INGREDIENTS:

ingredients from mineral origin, ingredients from vegetal origin, ingredients from marine origin, menthol.

**PACKAGE:** 25 kg paper bags



**DISTRIBUTED BY:** ANIMAXX-PREMIUM LTD.  
 2370 DABAS, OND VEZÉR ÚTJA 9.

**COUNTRY OF ORIGIN:** FRANCIAORSZÁG  
**TELEPHONE:** +36 30 792 64 17

**REGISTRATION NUMBER:**  
 766/1/2016 NÉBIH ÁTI (25 KG)

### MAIN ADVANTAGES:



Higher milk yield  
 Drop of clinical mastitis  
 Dry off hooves



Avoid humidity and skin problems  
 Dry off hooves



Avoid humidity and skin problems  
 Dry off hooves



Decrease mortality  
 recommended to avoid piglet mortality



Solve sole ulcer



Decrease of hoof disorders



#### POULTRY:

Turkey: 50 g per twice per week from the 2nd to 10th week.

Broiler: 50 g per twice per week from 2nd to 5th week.

Duck on straw: 70 g per m<sup>2</sup> every day from 1st to 12th week.

Ducks on duckboards: 50 g per m<sup>2</sup> every day from 1st to 12th week.



#### HORSES:

Before bedding renewing, spread 3 kg of Top Dry. Then in maintenance and before bedding aeration, spread 1 kg of Top Dry 2 to 3 times per week depending on the bedding moisture level.



#### SHEEP/GOATS:

65 g/m<sup>2</sup>/2x per week

AnimaXx Premium